



n-Butyric Acid AF

A HIGH QUALITY CARBOXYLIC ACID
FOR ANIMAL FEED



MATERIAL PROPERTIES

N-BUTYRIC ACID AF – A HIGH QUALITY PRODUCT FOR ANIMAL FEED

Supporting today's trend towards the replacement of antibiotic growth promoters by non-hazardous alternatives, Oxea offers carboxylic acids as special animal feed grades for use as raw material in agricultural industry. This trend started with the ban of antibiotic growth promoters in the European Union and expands to North America and Asia in recent years. Current African swine fever epidemic

puts additional pressure on farmers, not only in Asia, to raise animals even more efficiently.

OXEA's n-Butyric Acid AF – animal feed grade – is the key feedstock for different butyrate salts and glycerine esters – so-called butyrins. Salts are incorporated into feed premixes as fat-coated or non-coated administrative forms to channel the active ingredient to the

target organs in the animal's intestine. OXEA's n-Butyric Acid AF is produced and handled according to GMP (Good Manufacturing Practice) and HACCP (Hazard Analysis Critical Control Point).

Property	Limit	Unit	Test method
n-Butyric acid	min. 99.5	% (w/w)	gas chromatography
Propionic acid	max. 0.3	% (w/w)	gas chromatography
Isobutyric acid	max. 0.1	% (w/w)	gas chromatography
Water	max. 0.1	% (w/w)	DIN 51 777 / ASTM D 1364 (mod. Karl-Fischer-Method)
Platinum/Cobalt color (Hazen/APHA color)	max. 15		DIN EN 1557 / DIN ISO 6271 / ASTM D 1209

Figure 1: OXEA n-Butyric Acid AF (animal feed grade, CAS 107-92-6)

BUTYRIC ACID AS KEY RAW MATERIAL FOR NON-ANTIBIOTIC FEED ADDITIVES

The increasing awareness of the huge risk to human and animal health of the massive misuse of antibiotics as growth promoters in livestock led to a paradigm shift in Europe, North America, a growing number of countries in Asia and the rest of the world. For several years now, antibiotics

consumption on farms in Europe has decreased rapidly.

For example, in Germany, the overall sales of antibiotics used in livestock have been cut in more than half within a 2013 to 2017 timeframe. In other major animal producing countries,

such as the US, a similar trend started in the year 2016 by the voluntary action of big food companies. This led to a dramatic drop in antibiotic sales to farmers there.

Development of the sales of antimicrobials for farm animals in the US and in six key countries in Europe




Figure 2: Use of antimicrobials in six European countries (EU6 means: Germany, Spain, Netherlands, France, Denmark and Italy) and in the US. Sources: USDA, FIDIN, ESVAC report.

FEED ADDITIVES CAN COMPENSATE PRODUCTIVITY LOSS BY ANTIBIOTIC REDUCTION

In order to avoid the unfavorable effects seen by the removal of antibiotic growth promoters, farmers try to compensate their losses in livestock productivity by an intensified feeding of special feed additives like n-Butyric Acid derivatives.

Pathogen control, enteritis/diarrhea protection, and other health promoting measures to promote animal welfare in general are another important goal to be achieved by applying non-antibiotic feed additives.

Other effects like higher soy-protein contents in fish diet and improved rumen development in post-weaning calves prove the broad application range of n-Butyric Acid derivatives in many different farm animal species.

Less use of antibiotic growth promoters calls for rebalancing animals' health and performance




Figure 3: While the use of antibiotic growth promoters is declining, negative impact on performance and health of farm animals has to be compensated by feed additives like n-Butyric Acid derivatives, see also figure 2.

N-BUTYRIC ACID DERIVATIVES ADD VALUE IN ANIMAL HUSBANDRY BY ENHANCING ANIMALS' WELFARE AND PERFORMANCE

One of the most promising precursor molecules for feed additives is n-Butyric acid in its salt form or as mono-/tri-glyceride ester.

n-Butyric Acid is a natural substance present in biological liquids and tissues. It is naturally produced by the gastro-intestinal microbiota. Ruminants produce up to 1 kg butyrates each day while digesting fibers in the rumen. This endogenously provided breakdown product is directly used as an energy source by the animal's metabolism. Numerous scientific studies reveal that orally ingested n-Butyric Acid derivatives, like butyrates or butyrins*, exert several beneficial effects on different farmed animals. Researchers figured out that n-Butyric Acid derivatives can act directly – e.g. by controlling pathogens, by enlarging villi surfaces or by providing extra energy to enterocytes. But there is also an indirect mode of action e.g. by stimulating immune response. For a review see Bedford A et al, 2018 [29].

n-Butyric Acid derivatives show beneficial effects in poultry, aquaculture and cattle




Figure 4: Beneficial effect from using n-Butyric Acid derivatives as in-feed additive in different farm animal species. For references, see below.


SCIENTIFIC STUDIES EMPHASIZE EFFECTIVENESS OF N-BUTYRIC ACID DERIVATIVES AS A BENEFICIAL FEED ADDITIVE

Effectiveness of n-Butyric Acid derivatives is seen for in-feed application for farm animals like poultry, calves and different fish and shrimp species. Dozens of peer-reviewed scientific studies illustrate these beneficial effects when added to livestock diets.

Over the past decades feed conversion and growth rate in chicken production improved dramatically with the consequence that nutrition and health care gained much more relevance. In broiler and hen husbandry improved Salmonella control is one main focus in addition to general productivity improvements by feeding n-Butyric Acid derivatives.

Livestock		n-butyrate				
Poultry species	Animals tested in study	Sodium salt	Calcium salt	Fat coating	Feed conversion	Weight gain
Broiler	480	✓		✓	✓	
Broiler	200	✓		✓		✓
Laying hen	288		✓	✓	✓	
Laying hen	1000		✓			
Broiler	480	✓				
Broiler	96	✓		✓		✓
Broiler	2208	✓		✓		✓
Broiler	288	✓			✓	
Yellow broiler	180	✓		✓		
Broiler	150	✓		✓		
Breeder rooster	300	✓		✓		
Broiler	960		✓	✓	✓	✓
Broiler	720	✓		✓		
Broiler	240	✓			✓	✓

Productivity improvements				Health			Reference	
Epithelial surface	Egg quantity/quality	Reproduction	Feed digestibility	Pathogen protection	Intestinal health	Beneficial microbiota	Carcass state	
				✓		✓		Zou 2010 ¹
✓			✓					Smulikowska 2009 ²
	✓							Hamme 2013 ³
	✓							Sengor 2007 ⁴
				✓				Cerisuelo 2014 ⁵
✓								Czerwinski 2012 ⁶
						✓		Bortoluzzi 2017 ⁷
			✓					Qaisrani 2015 ⁸
					✓	✓		Zou 2019 ⁹
				✓				Fernández-Rubio 2009 ¹⁰
		✓						Alhaj 2018 ¹¹
✓			✓					Kaczmarek 2016 ¹²
✓						✓		Wu 2018 ¹³
✓				✓			✓	Panda 2009 ¹⁴



In aquaculture, farmed fish and shellfish are also beneficiaries of positive effects by being fed with n-Butyric Acid derivatives-containing diets.


Published scientific studies show that beside general productivity improvement by adding n-Butyric Acid derivatives, a higher acceptance of high-soy-protein-containing diets could be noted. This might open the door to more sustainable aquaculture by decreasing the use of fishmeal that runs short.

Livestock	n-butyrate	
Marine species	Animals tested in study	Sodium salt
Shrimp	3150	
Juvenile Sea Bream	120	
Juvenile Turbot	270	
Juvenile Gilthead Sea Bream	510	
Juvenile Gilthead Sea Bream	2200	
Juvenile common Carp	480	



Productivity improvements						Health		Reference	
Fat coating	Feed conversion	Increased weight gain	Epithelial surface	Soybean acceptance	Feed digestibility	Pathogen control	Lower mortality	Enteritis protection	
		✓			✓	✓	✓		da Silva 2014 ¹⁵
	✓	✓			✓				Robles 2013 ¹⁶
		✓	✓	✓				✓	Liu 2019 ¹⁷
				✓					Estensoro 2016 ¹⁸
				✓		✓		✓	Piazzon 2017 ¹⁹
	✓	✓	✓	✓				✓	Liu 2014 ²⁰





In recent years, cattle have also been tested for health and productivity improvements by diets and milk replacers being enriched with n-Butyric Acid derivatives. A growing number of studies prove beneficial effects like antibiotic

replacement, offer better feed conversion or improved health status, showing that ruminants can also profit from the advantages of different n-butyric acid administration forms.

Here the focus is clearly on calves with immature rumen where rapid development is needed to fulfill the economic need of an early weaning of calves from milk. The rumen passage in adult animals displays the key hurdle in applying n-Butyric Acid derivatives in beef and milk production.

Livestock	n-butyrate	Productivity improvements							Health			Reference		
		Butyric acid	Sodium salt	Fat coating	Feed conversion	Weight gain	Rumen development	Intestine maturation	Feed digestibility	Cell proliferation	Microbiota modulation		Less apoptosis	Overall health
Cattle species	Animals tested in study													
Suckling Holstein calves	29				✓	✓								Roh 2018 ²¹
Male Holstein calves	28						✓	✓		✓		✓		Górka 2014 ²²
Male Holstein calves	21											✓		Górka 2011a ²³
Male Holstein calves	28					✓			✓				✓	Górka 2011b ²⁴
Male Holstein calves	88				✓	✓							✓	Hill 2011 ²⁵
Holstein calves	88				✓	✓		✓						Guilloteau 2009 ²⁶
Holstein/Frisian calves	16				✓	✓					✓			O'Hara 2018 ²⁷
Male Holstein calves	48				✓	✓			✓					Hill 2016 ²⁸



DISCLAIMER

These scientific studies have not yet been reviewed for effectiveness nor has recommended dosage been established by the FDA.

- 1) Zou Y et al, 2010, "Effects of Sodium Butyrate on Gut pH value, Gut VFA and Cecal Microflora of Broilers, Department of Animal Science and Technology", Shandong Agricultural University, Tai-an, Shandong, P. R. China, 271000
- 2) Smulikowska S et al, 2009, "The effect of fat-coated organic acid salts and a feed enzyme on growth performance, nutrient utilization, microflora activity, and morphology of the small intestine in broiler chickens", *Journal of Animal and Feed Sciences* 18(3):478–89
- 3) Hamme V et al, 2013, "Effect of an additive based on coated calcium-butyrate on egg production and egg quality in laying hens", *Actes des 10èmes Journées de la Recherche Avicole et Palmipèdes à Foie Gras du 26 au 28 mars, 2013, Institut Technique de l'Aviculture, La Rochelle, France, 793–97*
- 4) Sengor E et al, 2007, "Short communication effects of short chain fatty acid (SCFA) supplementation on performance and egg characteristics of old breeder hens", *South African Journal of Animal Science* 37(3):158–63
- 5) Cerisuelo A et al, 2014, "The impact of a specific blend of essential oil components and sodium butyrate in feed on growth performance and Salmonella counts in experimentally challenged broilers", *Poultry Science* 93(3):599–606
- 6) Czerwinski J et al, 2012, "Effects of sodium butyrate and salinomycin upon intestinal microbiota, mucosal morphology and performance of broiler chickens", *Archives of animal nutrition* 66(2):102–16
- 7) Bortoluzzi C et al, 2017, "Sodium butyrate improved performance while modulating the cecal microbiota and regulating the expression of intestinal immune-related genes of broiler chickens", *Poultry Science* 96(11):3981–93
- 8) Qaisrani S et al, 2015, "Diet structure, butyric acid, and fermentable carbo-hydrates influence growth performance, gut morphology, and cecal fermentation characteristics in broilers", *Poultry Science* 94(9):2152–64
- 9) Zou X et al, 2019, "Effects of sodium butyrate on intestinal health and gut microbiota composition during intestinal inflammation progression in broilers", *Poultry Science* 98(10):4449–56
- 10) Fernández-Rubio C et al, 2009, "Butyric acid-based feed additives help protect broiler chickens from Salmonella Enteritidis infection", *Poultry Science* 88(5):943–48
- 11) Alhaj H et al, 2018, "Effects of dietary sodium butyrate on reproduction in adult breeder roosters", *Animal Reproduction Science* 196:111–19
- 12) Kaczmarek S et al, 2016, "Effect of different doses of coated butyric acid on growth performance and energy utilization in broilers", *Poultry Science* 95(4):851–59
- 13) Wu W et al, 2018, "Dietary sodium butyrate improves intestinal development and function by modulating the microbial community in broilers", *PLoS ONE* 13(5):e0197762
- 14) Panda AK et al, 2009, "Effect of Butyric Acid on Performance, Gastrointestinal Tract Health and Carcass Characteristics in Broiler Chickens", *Asian-Australasian Journal of Animal Sciences* 22(7):1026–31
- 15) da Silva BC, 2014, "Butyrate and propionate improve the growth performance of Litopenaeus vannamei", *Aquaculture Research* 47(2):612–23
- 16) Robles R et al, 2013, "Effect of partially protected butyrate used as feed additive on growth and intestinal metabolism in sea bream (*Sparus aurata*)", *Fish physiology and biochemistry* 39(6):1567–80
- 17) Liu Y et al, 2019, "Sodium butyrate supplementation in high-soybean meal diets for turbot (*Scophthalmus maximus* L.): Effects on inflammatory status, mucosal barriers and microbiota in the intestine", *Fish & Shellfish Immunology* 88(5):65–75
- 18) Estensoro I et al, 2016, "Dietary Butyrate Helps to Restore the Intestinal Status of a Marine Teleost (*Sparus aurata*) Fed Extreme Diets Low in Fish Meal and Fish Oil", *PLoS One* 11(11): e0166564
- 19) Piazzon NC et al, 2017, "Under control: how a dietary additive can restore the gut microbiome and proteomic profile, and improve disease resilience in a marine teleostean fish fed vegetable diets", *Microbiome* 5(1):164
- 20) Liu W et al, 2014, "Effects of dietary micro-encapsulated sodium butyrate on growth, intestinal mucosal morphology, immune response and adhesive bacteria in juvenile common carp (*Cyprinus carpio*) pre-fed with or without oxidised oil", *The British journal of nutrition* 112(1):15–29
- 21) Roh S et al, 2018, "Effects of butyrate supplementation in antibiotic-free milk replacer and starter on growth performance in suckling calves", *Animal Science Journal* 89(10):1486–91
- 22) Górka P et al, 2014, "Effect of method of delivery of sodium butyrate on maturation of the small intestine in newborn calves", *Journal of Dairy Science* 97(2):1026–35
- 23) Górka P et al, 2011a, "Is rumen development in newborn calves affected by different liquid feeds and small intestine development?", *Journal of Dairy Science* 94(6):3002–13
- 24) Górka P et al, 2011b, "Effect of method of delivery of sodium butyrate on rumen development in newborn calves", *Journal of Dairy Science* 94(11):5578–88
- 25) Ill TM et al, 2011, "Fatty acid intake alters growth and immunity in milk-fed calves", *Journal of Dairy Science* 94(8):3936–48
- 26) Guilloteau P et al, 2009, "Sodium-butyrate as a growth promoter in milk replacer formula for young calves", *Journal of Dairy Science* 92(3):1038–49
- 27) O'Hara O et al, 2018, "Effect of a butyrate-fortified milk replacer on gastrointestinal microbiota and products of fermentation in artificially reared dairy calves at weaning", *Scientific Reports* 8:14901
- 28) Hill TM et al, 2016, "Effect of milk replacer feeding rate and functional fatty acids on dairy calf performance and digestion of nutrients", *Journal of Dairy Science* 99(8):6352–61
- 29) Bedford A et al, 2018, "Implications of butyrate and its derivatives for gut health and animal production", *Animal Nutrition* 4:151–59

* Butyrin means tri-n-butyglyceride or mono-glyceride or mixtures thereof



For further information please contact your nearest
OXEA Sales Representative.

USA

Phone: +1 346 378 7300

Germany

Phone: +49 2173 9993-0

info@oxea-chemicals.com



[www.oxea-chemicals.com/en/
contact/sales-cr.html](http://www.oxea-chemicals.com/en/contact/sales-cr.html)

Disclaimer

For industrial use only. The information contained herein is accurate to the best of our knowledge. We do not suggest or guarantee that any hazards listed herein are the only ones which exist. OXEA makes no warranty of any kind, express or implied, concerning the safe use of this material in your process or in combination with other substances. The user has the sole responsibility to determine the suitability of the materials for any use and the manner of use contemplated. The user must meet all applicable safety and health standards.

Imprint

OXEA GmbH
Rheinpromenade 4a
40789 Monheim am Rhein
Germany

Phone: +49 2173 9993-0
info@oxea-chemicals.com

Trade Register
Amtsgericht Düsseldorf HRB 79958

VAT-Identification Number:
DE813758906

Layout

vE&K Advertising agency
Essen

www.oxea-chemicals.com